




Best Agrolife Limited

(An ISO 9001 : 2015 Certified Company)

ZODIO - A Unique 3-Way Action



Dose:
200 ml
Per Acre



Presentation :

1 Ltr, 500 ml, 200 ml & 100 ml



ZODIO (Azoxystrobin 18.2% w/w + Difenoconazole 11.4% w/w SC)




Mode of Action ►

Zodio is a broad spectrum fungicide having systemic activity against all major pathogens.

Recommendation ►

Zodio is recommended for the control of Anthracnose and Powdery mildew of Chilli, Early blight and Late blight of Tomato, Blast & Sheath blight of Paddy, Blight & Downy mildew of Maize, Rust & Powdery mildew of Wheat, Leaf spot & Grey mildew of Cotton, Red rot, Smut & Rust of Sugarcane, Purple blotch, Stemphylium blight & Downy mildew of Onion & Leaf blotch, Leaf spot & Rhizome rot of Turmeric.

Crop (S)	Common Name of Diseases	Dosage / HA		Dilution in water (Litre)
		a.i. (% conc. or ml/l)	Formulation	
Wheat	Rust & Powdery mildew	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500
Tomato	Early blight and Late blight	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500
Paddy	Blast & Sheath blight	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500
Maize	Blight & Downy mildew	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500
Chilli	Anthracnose and Powdery mildew	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500
Cotton	Leaf Spot & Grey mildew	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500
Sugarcane	Red rot, Smut & Rust	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	400
Onion	Purple blotch, Stemphylium blight & Downy mildew	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500
Turmeric	Leaf blotch, Leaf spot & Rhizome rot	0.03% or 0.3 g/l	0.1% or 1 ml/ litre water	500



Time of Application ►

Weather condition - Don't spray during windy and rainy weather; **Soil** - Spray when there is sufficient moisture in soil.

Give 1-2 sprays (depending on disease pressure, repeat the spray) with knapsack or any other suitable sprayer having hollow cone or any other suitable nozzle at the time when crops are infested with above mentioned diseases.