



Best Agrolife Limited

(An ISO 9001 : 2015 Certified Company)

Non Selective
Post-Emergence
Herbicide -
Inhibits Glutamine
Synthetase



Dose:
1-1.32Ltr
Per Acre

Presentation : 1Ltr & 500ml

BESTIE

(Glufosinate Ammonium 13.5% W/W SL)



Mode of Action ▶

Bestie is a broad-spectrum herbicide and acts against a wide range of annual and perennial broadleaf weeds and grasses. It is effective against a broad range of weeds, eliminating the need to apply several herbicides to control different weeds on a given crop. Its unique mode of action is a vital tool in weed resistance management, helping to improve crop health & agricultural yields. This unique mode of action makes it ideal to be used in rotation with other herbicides to mitigate weed resistance.

Recommendation ▶

Crop (S)	Common Name of Weeds	Dosage / HA		Dilution in water (Litre)
		ai (gm)	Formulation (Liter)	
Tea	Imperata cylindrica, Panicum repens, Borreria hispida, Digitaria sanguinalis, Commelina benghalensis, Ageratum conyzoides, Eleusine indica, Paspalum conjugatum.	375-500	2.5-3.3	375-500
Cotton	Echinochloa sp., Cynodon dactylan, Cyperus rotundus, Digitaria marginata, Dactyloctenium aegyptium	375-450	2.5-3.0	500

Method of use ▶

Spraying with knapsack sprayer with a flat fan or flood jet nozzle fitted with a spray shield.

Time of Application ▶

Cotton : Bestie is to be used as post emergence herbicide. It is to be applied when weeds are in active vegetative growth stage.

Tea : Bestie is to be used @ 375-500 gm a.i./ha. (2.5-3.3 lit/ha.) as post emergence herbicide. It is to be applied when weeds are at active vegetative growth stage/flowering stage with a spray shield to prevent drift on the tea plant.



BESTIE