



Best Agrolife Limited

(An ISO 9001 : 2015 Certified Company)


**Dual Mode Of Action -
Controls Multiple Stages
Of Fungus Development**



**Dose:
300 ml
Per Acre**

Presentation :

1 Ltr, 500 ml, 250 ml & 100 ml



ZOXIT

(Azoxystrobin 11% + Tebuconazole 18.3% w/w SC)




Mode of Action ▶

Zoxit is a suspensible concentrate fungicide for control of Fruit rot, Dieback and Powdery mildew in Chilli, Purple blotch (*Alternaria porri*) on Onion and Sheath blight on Rice Crop.

Recommendation ▶

Crop (S)	Common Name of Diseases	Dosage / HA		Dilution in water (Litre)
		a.i. (gms)	Formulation (ml)	
Rice	Sheath blight (<i>Rhizoctonia solani</i>)	82.5+137.25	750	800
Onion	Purple blotch (<i>Alternaria porri</i>)	82.5+137.25	750	800
Chilli	Fruit rot, Powdery mildew, Dieback	72+120	600	500-750
Potato	Early blight (<i>Alternaria solani</i>), Late blight (<i>Phytophthora infestans</i>)	82.5+137.25	750	500
Wheat	Yellow rust (<i>Puccinia striiformis</i>)	82.5+137.25	750	500
Tomato	Early blight (<i>Alternaria solani</i>)	82.5+137.25	750	500



Method of Use ▶

Take the measured quantity of fungicide and add to it a small quantity of water. Stir thoroughly using a stick or rod and then add remaining quantity of water, stir well to form final spray solution.

Time of Application ▶

To be applied preventively and to be repeated at 15 days interval depending upon prevailing weather conditions and disease pressure.